

# WEIGHTED COMPOSITION OPERATORS ON WEAK VECTOR-VALUED WEIGHTED BERGMAN SPACES AND HARDY SPACES

MOSTAFA HASSANLOU AND HAMID VAEZI

ABSTRACT. In this paper we investigate weighted composition operators between weak and strong vector-valued weighted Bergman spaces and Hardy spaces.

## 1. Introduction and Preliminaries

Weighted composition operators have been studied on different spaces of analytic functions. In [5], Contreras and Hernandez-Diaz have made a study of weighted composition operators on Hardy spaces whereas Mirzakarimi and Siddighi [12] have studied these operators on Bergman and Dirichlet spaces. On Bloch-type spaces, these operators are explored by MacCluer and Zhao [11], Ohno [13], Ohno and Zhao [14] and Ohno, Stroethoff and Zhao [15]. In [8] Kumar studied weighted composition operators between spaces of Dirichlet type by using Carleson measures.

Recently these studies are about spaces of vector-valued analytic functions. For example, in [17], Wang presented some necessary and sufficient conditions for weighted composition operators to be bounded on vector-valued Dirichlet spaces and Laitila, Tylli and Wang [10] studied composition operators from weak to strong vector-valued Bergman spaces Hardy spaces. For some information about vector-valued Bergman spaces see [1, 3].

Let  $X$  be a complex Banach space and  $\mathbb{D}$  be the open unit ball of  $\mathbb{C}$ . We consider *weight* as a strictly positive bounded continuous function  $v : \mathbb{D} \rightarrow \mathbb{R}^+$ . Let  $p \geq 1$  and  $v$  be a weight. The vector-valued weighted Bergman space  $A_v^p(X)$  consists of all analytic functions  $f : \mathbb{D} \rightarrow X$  such that

$$\|f\|_{A_v^p(X)} = \left( \int_{\mathbb{D}} \|f(z)\|_X^p v(z) dA(z) \right)^{\frac{1}{p}} < \infty.$$

where  $dA$  is the normalized area measure on  $\mathbb{D}$ . Also, the vector-valued weighted Hardy space  $H_v^p(X)$  consists of all analytic functions  $f : \mathbb{D} \rightarrow X$  for which

$$\|f\|_{H_v^p(X)} = \sup_{0 < r < 1} \left( \int_{\mathbb{T}} \|f(r\zeta)\|_X^p v(r\zeta) dm(\zeta) \right)^{\frac{1}{p}} < \infty,$$

where  $dm(\zeta)$  is the normalized Lebesgue measure on the unit circle  $\mathbb{T} = \partial\mathbb{D}$ . In the case  $X = \mathbb{C}$ , we write  $A_v^p(X) = A_v^p$  and  $H_v^p(X) = H_v^p$ . Also, if  $v \equiv 1$ , then we have  $A_v^p(X) = A^p(X)$  and  $H_v^p(X) = H^p(X)$ . The following weak versions of these spaces were considered by e.g. Blasco [2] and Bonet, Domanski and Lindstrom [4]:

---

2000 *Mathematics Subject Classification.* 47B35.

*Key words and phrases.* Weighted composition operators, vector-valued Bergman spaces, vector-valued Hardy spaces .

the weak spaces  $wA_v^p(X)$  and  $wH_v^p(X)$  consist of all analytic functions  $f : \mathbb{D} \rightarrow X$  for which

$$\|f\|_{wA_v^p(X)} = \sup_{\|x^*\| \leq 1} \|x^* \circ f\|_{A_v^p}, \quad \|f\|_{wH_v^p(X)} = \sup_{\|x^*\| \leq 1} \|x^* \circ f\|_{H_v^p},$$

are finite, respectively. Here  $x^* \in X^*$ , the dual space of  $X$ .

Let  $\varphi$  be an analytic self-map of  $\mathbb{D}$ ; that is  $\varphi(\mathbb{D}) \subset \mathbb{D}$ , and  $u$  a scalar-valued analytic function on  $\mathbb{D}$ . We can define the weighted composition operator  $uC_\varphi$  on the space of analytic functions as follows:

$$uC_\varphi(f)(z) = u(z)f(\varphi(z)).$$

When  $u(z) \equiv 1$ , we just have the composition operator  $C_\varphi$ , defined by  $C_\varphi(f) = f \circ \varphi$ . Also if  $\varphi = I$ , the identity function, then we get the multiplication operator  $M_u$  defined by  $M_u(f)(z) = u(z)f(z)$ . It is well known that for every analytic map  $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ ,  $C_\varphi : A^p(X) \rightarrow A^p(X)$  and  $C_\varphi : H^p(X) \rightarrow H^p(X)$  are bounded, and also on  $wA^p(X), wH^p(X)$ . For complete discussion about composition operators we refer to [6, 16]. We consider the infinite dimensional complex Banach space  $X$ , since  $wA^p(X) = A^p(X)$  and  $wH^p(X) = H^p(X)$ , for  $\alpha > -1$  and any finite dimensional Banach space  $X$ .

But for the infinite dimensional complex Banach space  $X$ ,  $A^p(X) \neq wA^p(X)$  ( $H^p(X) \neq wH^p(X)$ ) and  $\|\cdot\|_{wA^p(X)}$  is not equivalent to  $\|\cdot\|_{A^p(X)}$  on  $A^p(X)$  ( $\|\cdot\|_{wH^p(X)}$  is not equivalent to  $\|\cdot\|_{H^p(X)}$  on  $H^p(X)$ ), see [10] Proposition 3.1 ([9] Example 15).

Our aim in this paper is to compute the norm of weighted composition operators between  $wA_v^p(X)$  and  $A_{v'}^p(X)$ , for  $p \geq 2$  and also between  $wH_v^p(X)$  and  $H_{v'}^p(X)$ , for  $p \geq 2$ , where  $v$  and  $v'$  are weights.

Throughout the remainder of this paper,  $c$  will denote a positive constant, the exact value of which will vary from one appearance to the next. The notation  $A \approx B$  means that there are positive constants  $c_1$  and  $c_2$  such that  $c_1 A \leq B \leq c_2 A$ .

## 2. Main Results

**Proposition 2.1.** Let  $X$  be any complex Banach spaces,  $v$  be a weight of the form  $v = |\mu|$ , where  $\mu$  is an analytic function without any zeros on  $\mathbb{D}$ ,  $v'$  be a weight and  $1 \leq p < \infty$ . Then

$$(2.1) \quad \|uC_\varphi : wA_v^p(X) \longrightarrow A_{v'}^p(X)\| \leq \left( \int_{\mathbb{D}} \frac{|u(z)|^p v'(z)}{(1 - |\varphi(z)|^2)^2 v(\varphi(z))} dA(z) \right)^{1/p}, \text{ and}$$

$$(2.2) \quad \|uC_\varphi : wH_v^p(X) \longrightarrow H_{v'}^p(X)\| \leq \sup_{0 < r < 1} \left( \int_{\mathbb{T}} \frac{|u(r\zeta)|^p v'(r\zeta)}{(1 - |\varphi(r\zeta)|^2)^2 v(\varphi(r\zeta))} dm(\zeta) \right)^{1/p}.$$

**Proof.** By Lemma 2.1 of [18] we have

$$|f(z)| \leq \frac{\|f\|_{A_v^p}}{(1 - |z|^2)^{\frac{2}{p}} v(z)^{\frac{1}{p}}},$$

for any  $f \in A_v^p$  and  $z \in \mathbb{D}$ . Thus, for  $f \in wA_v^p(X)$ , we have

$$\|f(z)\|_X^p = \sup_{\|x^*\| \leq 1} |(x^* \circ f)(z)|^p \leq \frac{1}{(1 - |z|^2)^2 v(z)} \sup_{\|x^*\| \leq 1} \|x^* \circ f\|_{A_v^p}^p$$

$$= \frac{1}{(1 - |z|^2)^2 v(z)} \|f\|_{wA_v^p(X)}^p.$$

Hence

$$\begin{aligned} \|uC_\varphi f\|_{A_{v'}^p(X)}^p &= \int_{\mathbb{D}} |u(z)|^p \|f(\varphi(z))\|_X^p v'(z) dA(z) \\ &\leq \|f\|_{wA_v^p(X)}^p \int_{\mathbb{D}} \frac{|u(z)|^p v'(z)}{(1 - |z|^2)^2 v(\varphi(z))} dA(z). \end{aligned}$$

The proof of the theorem is complete.  $\square$

For the next results we need the following Dvoretzky's well-known theorem.

**Lemma 2.2.** [7] Suppose that  $X$  is an infinite dimensional complex Banach space. Then for any  $\epsilon > 0$  and  $n \in \mathbb{N}$ , there is a linear embedding  $T_n : l_n^2 \rightarrow X$  such that

$$(2.3) \quad (1 + \epsilon)^{-1} \left( \sum_{j=1}^n |a_j|^2 \right)^{1/2} \leq \left\| \sum_{j=1}^n a_j T_n e_j \right\|_X \leq \left( \sum_{j=1}^n |a_j|^2 \right)^{1/2}$$

for any scalars  $a_1, a_2, \dots, a_n$  and some orthonormal basis  $\{e_1, \dots, e_n\}$  of  $l_n^2$ .

Now, we prove a lower bound for the operator  $uC_\varphi : wA_v^p(X) \rightarrow A_{v'}^p(X)$ , in the case  $2 \leq p < \infty$ .

**Theorem 2.3.** Let  $X$  be any complex infinite-dimensional Banach space,  $v$  be a weight of the form  $v = |\mu|$ , where  $\mu$  is an analytic function without any zeros on  $\mathbb{D}$ ,  $v'$  be a weight and  $2 \leq p < \infty$ . Then

$$(2.4) \quad \|uC_\varphi : wA_v^p(X) \rightarrow A_{v'}^p(X)\| \approx \left( \int_{\mathbb{D}} \frac{|u(z)|^p v'(z)}{(1 - |\varphi(z)|^2)^2 v(\varphi(z))} dA(z) \right)^{1/p}.$$

**Proof.** We only prove there exists a positive constant  $c$  such that

$$\|uC_\varphi : wA_v^p(X) \rightarrow A_{v'}^p(X)\| \geq c \left( \int_{\mathbb{D}} \frac{|u(z)|^p v'(z)}{(1 - |\varphi(z)|^2)^2 v(\varphi(z))} dA(z) \right)^{1/p}.$$

Suppose that  $x \in X$  with  $\|x\| = 1$  and define  $g : \mathbb{D} \rightarrow X$  by  $g(z) = \frac{1}{\mu(z)^{\frac{1}{p}}} x$ . Then  $g$  is an analytic function on  $\mathbb{D}$ , and  $\|g\|_{wA_v^p(X)} = 1$ , so that

$$\|uC_\varphi\|^p \geq \|ug \circ \varphi\|_{A_{v'}^p}^p = \int_{\mathbb{D}} \frac{|u(z)|^p v'(z)}{v(\varphi(z))} dA(z).$$

Hence

$$\int_{\{z \in \mathbb{D} : |\varphi(z)|^2 < 1/2\}} \frac{|u(z)|^p v'(z)}{(1 - |\varphi(z)|^2)^2 v(\varphi(z))} dA(z) \leq 4 \int_{\mathbb{D}} \frac{|u(z)|^p v'(z)}{v(\varphi(z))} dA(z) \leq 4 \|uC_\varphi\|^p.$$

So, it will be sufficient to prove that there exists a positive constant  $c$  such that

$$\|uC_\varphi\|^p \geq c \int_{\{z \in \mathbb{D} : |\varphi(z)|^2 \geq 1/2\}} \frac{|u(z)|^p v'(z)}{(1 - |\varphi(z)|^2)^2 v(\varphi(z))} dA(z).$$

Let  $\lambda_k = k^{2/p-1/2}$ , for any  $n \in \mathbb{N}$ , we define functions  $f_n$  as follows

$$f_n(z) = \frac{1}{\mu(z)^{\frac{1}{p}}} \sum_{k=1}^n \lambda_k z^k T_n e_k,$$

where the linear embedding  $T_n$  is the same as in Lemma 2.2,  $\|T_n\| = 1$  and  $\|T_n^{-1}\| \leq (1 + \epsilon)$  and  $(e_1, \dots, e_n)$  is an orthonormal basis of  $\ell_2^n$ . As in the proof of Theorem 3.2 [10], there exists  $c > 0$  such that for  $X^*$  with  $\|x^*\| \leq 1$ , we have

$$\begin{aligned} \|x^* \circ f_n\|_{A_v^p} &= \left\| \frac{1}{\mu(z)^{\frac{1}{p}}} \sum_{k=1}^n \lambda_k z^k x^* T_n e_k \right\|_{A_v^p} \\ &= \left\| \sum_{k=1}^n \lambda_k x^* (T_n e_k) z^k \right\|_{A^p} \\ &\leq c \left( \sum_{k=1}^n |x^* (T_n e_k)|^2 \right)^{1/2} \leq c. \end{aligned}$$

It follows that  $\|f_n\|_{wA_v^p(X)} \leq c$ . Thus, Fatou's lemma implies that

$$\begin{aligned} \|uC_\varphi\|^p &\geq c^{-p} \limsup_{n \rightarrow \infty} \|uC_\varphi f_n\|_{A_{v'}^p(X)}^p \\ &= c^{-p} \limsup_{n \rightarrow \infty} \int_{\mathbb{D}} |u(z)|^p \left\| \frac{1}{\mu(\varphi(z))^{\frac{1}{p}}} \sum_{k=1}^n \lambda_k \varphi(z)^k T_n e_k \right\|_X^p v'(z) dA(z) \\ &= c^{-p} \limsup_{n \rightarrow \infty} \int_{\mathbb{D}} \left\| \sum_{k=1}^n \lambda_k \varphi(z)^k T_n e_k \right\|_X^p \frac{|u(z)|^p v'(z)}{v(\varphi(z))} dA(z) \\ &\geq \frac{c^{-p}}{(1 + \epsilon)^p} \limsup_{n \rightarrow \infty} \int_{\mathbb{D}} \left( \sum_{k=1}^n k^{4/p-1} |\varphi(z)|^{2k} \right)^{p/2} \frac{|u(z)|^p v'(z)}{v(\varphi(z))} dA(z) \\ &= \frac{c^{-p}}{(1 + \epsilon)^p} \int_{\mathbb{D}} \left( \sum_{k=1}^{\infty} k^{4/p-1} |\varphi(z)|^{2k} \right)^{p/2} \frac{|u(z)|^p v'(z)}{v(\varphi(z))} dA(z) \\ &\geq \frac{c_1 c^{-p}}{(1 + \epsilon)^p} \int_{\{z \in \mathbb{D} : |\varphi(z)|^2 \geq 1/2\}} \frac{|u(z)|^p v'(z)}{(1 - |\varphi(z)|^2)^2 v(\varphi(z))} dA(z) \end{aligned}$$

and the last inequality is derived by Lemma 2.3 [10]. As  $\epsilon > 0$  was arbitrary, we obtain the desired lower bound estimate.

□

**Theorem 2.4.** Let  $X$  be any complex infinite-dimensional Banach space,  $v$  be a weight of the form  $v = |\mu|$ , where  $\mu$  is an analytic function without any zeros on  $\mathbb{D}$ ,  $v'$  be a weight and  $2 \leq p < \infty$ . Then

$$(2.5) \quad \|uC_\varphi : wH_v^p(X) \longrightarrow H_{v'}^p(X)\| \approx \left( \int_{\mathbb{T}} \frac{|u(\zeta)|^p v'(\zeta)}{(1 - |\varphi(\zeta)|^2) v(\varphi(\zeta))} dm(\zeta) \right)^{1/p}.$$

**Proof.** Similar to the proof of previous theorem, we only prove that there exists  $c > 0$  such that

$$\|uC_\varphi\|^p \geq c \int_{\{\zeta \in \mathbb{T} : |\varphi(r\zeta)|^2 \geq 1/2\}} \frac{|u(r\zeta)|^p v'(r\zeta)}{(1 - |\varphi(r\zeta)|^2) v(\varphi(r\zeta))} dm(\zeta).$$

Let  $\lambda_k = k^{1/p-1/2}$  and define

$$f_n(z) := \frac{1}{\mu(z)^{\frac{1}{p}}} \sum_{k=1}^n \lambda_k z^k T_n e_k,$$

where the linear embedding  $T_n$  is the same as in Lemma 2.2,  $\|T_n\| = 1$  and  $\|T_n^{-1}\| \leq (1 + \epsilon)$  and  $(e_1, \dots, e_n)$  is an orthonormal basis of  $\ell_2^n$ . As in the proof of Theorem 2.2 [10], there exists  $c > 0$  such that for  $X^*$  with  $\|x^*\| \leq 1$ , we have

$$\begin{aligned} \|x^* \circ f_n\|_{H_v^p} &= \left\| \frac{1}{\mu(z)^{\frac{1}{p}}} \sum_{k=1}^n \lambda_k z^k x^* T_n e_k \right\|_{H_v^p} \\ &= \left\| \sum_{k=1}^n \lambda_k x^* (T_n e_k) z^k \right\|_{H^p} \\ &\leq c \left( \sum_{k=1}^n |x^* (T_n e_k)|^2 \right)^{1/2} \leq c. \end{aligned}$$

Thus  $\|f_n\|_{wH_v^p(X)} \leq c$  and by using Fatou's lemma and Lemma 2.3 [10], we have

$$\begin{aligned} \|uC_\varphi\|^p &\geq c^{-p} \limsup_{n \rightarrow \infty} \|uC_\varphi f_n\|_{H_v^p(X)}^p \\ &= c^{-p} \limsup_{n \rightarrow \infty} \int_{\mathbb{T}} |u(r\zeta)|^p \left\| \frac{1}{\mu(\varphi(r\zeta))^{\frac{1}{p}}} \sum_{k=1}^n \lambda_k \varphi(r\zeta)^k T_n e_k \right\|_X^p v'(r\zeta) dm(\zeta) \\ &= c^{-p} \limsup_{n \rightarrow \infty} \int_{\mathbb{T}} \left\| \sum_{k=1}^n \lambda_k \varphi(r\zeta)^k T_n e_k \right\|_X^p \frac{|u(r\zeta)|^p v'(r\zeta)}{v(\varphi(r\zeta))} dm(\zeta) \\ &\geq \frac{c^{-p}}{(1 + \epsilon)^p} \limsup_{n \rightarrow \infty} \int_{\mathbb{T}} \left( \sum_{k=1}^n k^{2/p-1} |\varphi(r\zeta)|^{2k} \right)^{p/2} \frac{|u(r\zeta)|^p v'(r\zeta)}{v(\varphi(r\zeta))} dm(\zeta) \\ &= \frac{c^{-p}}{(1 + \epsilon)^p} \int_{\mathbb{D}} \left( \sum_{k=1}^{\infty} k^{2/p-1} |\varphi(r\zeta)|^{2k} \right)^{p/2} \frac{|u(r\zeta)|^p v'(r\zeta)}{v(\varphi(r\zeta))} dm(\zeta) \\ &\geq \frac{c_1 c^{-p}}{(1 + \epsilon)^p} \int_{\{z \in \mathbb{T} : |\varphi(r\zeta)|^2 \geq 1/2\}} \frac{|u(r\zeta)|^p v'(r\zeta)}{(1 - |\varphi(r\zeta)|^2) v(\varphi(r\zeta))} dm(\zeta). \end{aligned}$$

Take  $\epsilon = 1$ , then

$$\|uC_\varphi\|^p \geq c \int_{\mathbb{T}} \frac{|u(r\zeta)|^p v'(r\zeta)}{(1 - |\varphi(r\zeta)|^2) v(\varphi(r\zeta))} dm(\zeta).$$

As  $r \rightarrow 1$ ,

$$\begin{aligned} \|uC_\varphi\|^p &\geq c \limsup_{r \rightarrow 1} \int_{\mathbb{T}} \frac{|u(r\zeta)|^p v'(r\zeta)}{(1 - |\varphi(r\zeta)|^2) v(\varphi(r\zeta))} dm(\zeta) \\ &\geq c \int_{\mathbb{T}} \frac{|u(\zeta)|^p v'(\zeta)}{(1 - |\varphi(\zeta)|^2) v(\varphi(\zeta))} dm(\zeta). \end{aligned}$$

**Acknowledgment.**

#### REFERENCES

- [1] J. ARREGUI, O. BLASCO, *Bergman and Bloch spaces of vector-valued functions*, Math Nachr, 2003, 261/262: 3-22.
- [2] O. Blasco, *Boundary values of vector-valued harmonic functions considered as operators*, Studia Math. **86** (1987), 19-33.
- [3] O. BLASCO, *Introduction to vector valued Bergman spaces*, in *Function Spaces and Operator Theory* (Joensuu, 2003), Univ. Joensuu Dept. Math. Rep. Ser., vol. 8, Univ. Joensuu, Joensuu 2005, pp. 930.

- [4] J. Bonet, P. Domanski, M. Lindstrom, *Weakly compact composition operators on analytic vector-valued function spaces*, Ann. Acad. Sci. Fenn. Math. **26** (2001), 233-248.
- [5] M. D. CONTRERAS AND A. G. HERNANDEZ-DIAZ, *Weighted composition operators on Hardy spaces*, J. Math. Anal. Appl. 263 (2001), no. 1, 224-233.
- [6] C. C. COWEN AND B. D. MACCLUER, *Composition operators on spaces of analytic functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla, USA, 1995.
- [7] J. DIESTEL, H. JARCHOW, AND A. TONGE, *Absolutely Summing Operators*, Cambridge Univ. Press, Cambridge 1995.
- [8] S. KUMAR, *Weighted composition operators between spaces of Dirichlet type*, Rev Mat Complut, 2009, 22: 469-488.
- [9] J. Laitila, *Weakly compact composition operators on vector-valued BMOA*. J. Math. Anal. Appl., **308** (2005), 730-745.
- [10] J. LAITILA , H.-O. TYLLI, AND M. WANG , *Composition operators from weak to strong spaces of vector-valued analytic functions*, J Operator Theory, 2009, 62: 281-295
- [11] B. D. MACCLUER AND R. ZHAO, *Essential norms of weighted composition operators between Bloch-type spaces*, Rocky Mountain J. Math. 33 (2003), no. 4, 1437-1458.
- [12] G. MIRZAKARIMI AND K. SIDDIGHI, *Weighted composition operators on Bergman and Dirichlet spaces*, Georgian Math. J. 4 (1997), no. 4, 373-383.
- [13] S. OHNO, *Weighted composition operators between  $H^\infty$  and the Bloch space*, Taiwanese J. Math. 5 (2001), no. 3, 555-563.
- [14] S. OHNO AND R. ZHAO, *Weighted composition operators on the Bloch space*, Bull. Austral. Math. Soc. 63 (2001), no. 2, 177-185.
- [15] S. OHNO, K. STROETHOFF, AND R. ZHAO, *Weighted composition operators between Bloch-type spaces*, Rocky Mountain J. Math. 33 (2003), no. 1, 191-215.
- [16] J. H. SHAPIRO, *Composition operator and classical function theory*, Springer-Verlag, New York, 1993.
- [17] M. WANG, *Weighted composition operators between Dirichlet spaces*, Acta Math. Sci. 2011,31B(2):641-651.
- [18] E. WOLF, *Weighted composition operators between weighted Bergman spaces and weighted Banach spaces of holomorphic functions*, Rev. Mat. Complut. **21** (2008), 4267-4273.
- [19] K. ZHU, *Spaces of holomorphic functions in the unit ball*, Springer, New York, 2005.